

**ASSESSMENT AND MAPPING OF SOME IMPORTANT SOIL  
PARAMETERS INCLUDING SOIL ACIDITY FOR THE STATE OF  
JHARKHAND (1:50,000 SCALE) TOWARDS  
RATIONAL LAND USE PLAN**

**SIMDEGA DISTRICT**



*National Bureau of Soil Survey and Land Use Planning (ICAR)  
Regional Centre, Kolkata*

*In collaboration with :  
Deptt. Of Soil Science & Agricultural Chemistry, BAU, Ranchi, Jharkhand*

*Sponsored by : Department of Agriculture & Cane Development,  
Govt. of Jharkhand*

## **1. INTRODUCTION**

Reliable information on the location, extent and quality of soil and land resources is the first requirement in planning for the sustainable management of land resources. The components of land i.e., soils, climate, water, nutrient and biota are organised into eco-system which provide a variety of services that are essential to the maintenance of the life support system and the productive capacity of the environment. Our land mass is fixed, but the competition among different kinds of uses for this land is increasing because of rapidly rising global population. Therefore, integrated land resource planning and management are required to resolve these conflicts and soil resource survey seems to be a viable means in this process and knowledge of soil fertility status and problems of soils like soil acidity/alkalinity become essential for sustainable land use plan.

Soil fertility is an aspect of the soil-plant relationship. Fertility status of the soils is primarily and importantly dependent upon both the macro and micronutrient reserve of that soil. Continued removal of nutrients by crops, with little or no replacement will increase the nutrient stress in plants and ultimately lowers the productivity. The fertility status of the soils mainly depends on the nature of vegetation, climate, topography, texture of soil and decomposition rate of organic matter. Optimum productivity of any cropping systems depends on adequate supply of plant nutrients. GIS is a versatile tool used for integration of soil database and production of a variety of users specific and user-friendly interpretative maps. This further leads to accurately and scientifically interpret and plan some of the aspects like conservation of organic matter, soil reaction (pH) control and fertilization.

Keeping in view NBSS & LUP, Regional Centre, Kolkata in collaboration with Department of Soil Science and Agricultural Chemistry, BAU, Ranchi, Jharkhand undertook a project entitled "Assessment and mapping of some important soil parameters including soil acidity for the state of Jharkhand (1:50,000 scale)

towards rational land use plan” from Department of Agriculture, Govt. of Jharkhand. The major objectives of the project were

- Preparation of districtwise soil acidity maps
- Preparation of districtwise soil fertility maps (Organic carbon, available N, P, K, S and available Fe, Mn, Zn, Cu and B)

The above maps will provide information regarding soil nutrients and soil acidity status for the districts, which will be very useful in identification of site specific problems for planning purposes. The present report deals with the above mentioned objectives of the Simdega district, Jharkhand.

## **2. GENERAL DESCRIPTION OF THE AREA**

### **2.1 Location and Extent**

Simdega came into existence as a district on 30<sup>th</sup> April 2001. Prior to this it formed part of Gumla district as a subdivision. The district is divided into seven development blocks namely Bano, Jaldega, Kurdeg, Bolwa, Simdega, Kolebira and Thethaitangar. It is bounded by Gumla district in north, Chhatisgarh and Orissa state in the west and South respectively. The total geographical area of the district is 3757 sq. km and population is 5,47,969 persons (Census of India, 2001).

### **2.2 Physiography, Geology and Drainage**

The district consist of Simdega plateau with rugged topography with turbulent streams, steep slopes, high cliffs and narrow valleys. The general slope of the district is from north to South. Geologically the area is comprised with Archean granites and gneisses. In the uplands considerable thickness of laterite of Pleistocene age is found in the granite and gneisses tracts. Alluvium of recent to sub-recent age is found in the river valley. The area is drained by South Koel and Sankh river.

### **2.3 Climate**

Average annual rainfall of the area is about 900 mm. But in areas of high elevations, the amount of rainfall increases. The average temperature ranges between 16 to 29<sup>o</sup> C.

### **2.4 Agriculture and Land Use**

As the area is highly dissected and has rough terrain, contiguous agricultural lands are found in limited areas. The low lying *Don* area provide ideal condition for rice production and comparatively higher *Tanr* areas are grown for

ragi, maize, pulses and vegetables. There are smaller irrigation scheme in the form of reservoir.

### Land Use in Simdega District (1997-98)

	Simdega	Jharkhand
1. Forest	15.06 %	29.2 %
2. Net sown area	29.02 %	22.7 %
3. Barren and unculturable waste	7.37 %	7.2 %
4. Non agricultural use	6.95 %	9.9 %
5. Orchards	1.75 %	2.5 %
6. Pasture	0.05 %	
7. Culturable wasteland	5.07 %	3.5 %
8. Current and other fallow	34.73 %	25.0 %

Source: Fertilizer and Agriculture Statistics, Eastern Region (2003-2004)

## 2.5 Soils

The soils occurring in different landforms have been characterised during soil resource mapping of the state on 1:250,000 scale (Halder *et al.* 1996) and three soil orders namely Entisols, Inceptisols and Alfisols were observed in Simdega district (Fig.1 and table 1). Inceptisols covers 37.8 percent of total geographical area followed by Entisols (34.2 %) and Alfisols (27.0%)

**Table 1. Soils of the district and their extent**

Map unit	Taxonomy	Area ('00ha)	% of the TGA
15	Loamy-skeletal, mixed, hyperthermic Lithic Ustorthents Fine loamy, mixed, hyperthermic Ultic Haplustalfs	280	7.45
17	Loamy, mixed, hyperthermic Lithic Ustorthents Fine, mixed, hyperthermic Typic Rhodustalfs	12	0.32
18	Loamy, mixed, hyperthermic Lithic Ustorthents Fine loamy, mixed, hyperthermic Typic Haplustalfs	13	0.35
19	Loamy-skeletal, mixed hyperthermic Lithic Ustorthents Fine loamy, mixed, hyperthermic Typic Haplustepts	15	0.40
20	Loamy, mixed, hyperthermic Lithic Ustorthents Fine, mixed, hyperthermic Typic Rhodustalfs	50	1.33

<b>Map unit</b>	<b>Taxonomy</b>	<b>Area ('00ha)</b>	<b>% of the TGA</b>
21	Coarse loamy, mixed, hyperthermic Typic Ustorthents Fine, mixed, hyperthermic Rhodic Paleustalfs	115	3.06
22	Fine, mixed, hyperthermic Typic Paleustalfs Fine, mixed, hyperthermic Typic Rhodustalfs	25	0.66
24	Fine, mixed, hyperthermic Typic Haplustalfs Fine-loamy, mixed, hyperthermic Typic Haplustepts	81	2.16
35	Loamy-skeletal, mixed, hyperthermic Lithic Ustorthents Fine-loamy, mixed, hyperthermic Typic Haplustalfs	190	5.06
38	Fine loamy, mixed, hyperthermic Typic Paleustalfs Fine loamy, mixed, hyperthermic Typic Haplustepts	311	8.28
39	Fine, mixed, hyperthermic Rhodic Paleustalfs Fine-loamy, mixed, hyperthermic Typic Haplustepts	106	2.82
40	Fine loamy, mixed, hyperthermic Typic Haplustepts Fine loamy, mixed, hyperthermic Typic Haplustalfs	54	1.44
42	Fine, mixed, hyperthermic Typic Rhodustalfs Fine loamy, mixed, hyperthermic Typic Ustorthents	99	2.64
44	Fine, mixed, hyperthermic Aerico Endoaquepts Fine, mixed, hyperthermic Typic Haplustepts	31	0.83
50	Loamy, mixed, hyperthermic Lithic Ustorthents Fine loamy, mixed, hyperthermic Typic Haplustepts	534	14.21
51	Fine loamy, mixed, hyperthermic Typic Haplustepts Loamy, mixed, hyperthermic Lithic Ustorthents	29	0.77
53	Fine loamy, mixed, hyperthermic Typic Haplustepts Fine, mixed, hyperthermic Typic Haplustalfs	18	0.48
54	Loamy, mixed, hyperthermic Lithic Ustorthents Fine, mixed, hyperthermic Typic Haplustalfs	111	2.95
60	Fine loamy, mixed, hyperthermic Typic Haplustepts Fine, mixed, hyperthermic Typic Paleustalfs	128	3.41
62	Fine, mixed, hyperthermic Aerico Endoaquepts Fine loamy, mixed, hyperthermic Typic Haplustepts	61	1.62
64	Loamy, mixed, hyperthermic Lithic Ustorthents Fine, mixed, hyperthermic Typic Paleustalfs	144	3.83
65	Loamy, mixed, hyperthermic Lithic Ustorthents Fine loamy, mixed, hyperthermic Typic Haplustepts	298	7.93
66	Loamy-skeletal, mixed, hyperthermic Typic Haplustepts Coarse loamy, mixed, hyperthermic Typic Ustorthents	42	1.12
67	Coarse loamy, mixed, hyperthermic Typic Ustorthents Fine, mixed, hyperthermic Typic Haplustalfs	263	7.00
69	Fine loamy, mixed, hyperthermic Typic Haplustepts Fine loamy, mixed, hyperthermic Aerico Endoaquepts	67	1.78
70	Fine loamy, mixed, hyperthermic Typic Haplustepts Fine, mixed, hyperthermic Aerico Endoaquepts	112	2.98
71	Fine, mixed, hyperthermic Aerico Endoaquepts Fine, mixed, hyperthermic Typic Haplustepts	205	5.46
73	Fine, mixed, hyperthermic Aerico Endoaquepts Very fine, mixed, hyperthermic Vertic Endoaquepts	290	7.72
76	Fine loamy, mixed, hyperthermic Typic Haplustalfs Fine loamy, mixed, hyperthermic Typic Paleustalfs	35	0.93
Miscellaneous		38	1.01
Total		3757	100.00

### 3. METHODOLOGY

The base map of the district was prepared on 1:50,000 scale using Survey of India toposheets (73B/2,3,5,6,7,9,10,11,13,14,15 and 73F/1,2,3) and all the maps were demarcated with grid points at 2.5 km interval.

Surface soil samples from demarcated grid points and other related informations were collected through field survey. Soil samples were air dried, processed and analysed for pH, organic carbon, available phosphorous and potassium (Page *et al.*, 1982), available nitrogen (Subbaiah and Asija, 1956), available sulphur by using 0.15 percent  $\text{CaCl}_2$  as the extractant (William and Steinbergs, 1959), available (DTPA extractable) Fe, Mn, Zn and Cu (Lindsay and Norvell, 1978) and available B (hot water soluble) by Carmine method (Hatcher and Wilcox, 1950).

The soils are grouped under different soil reaction classes viz extremely acidic (pH<4.5), very strongly acidic (pH 4.5 – 5.0 ), strongly acidic (pH 5.1 – 5.5), moderately acidic (pH 5.6-6.0), slightly acidic (pH 6.1-6.5), neutral (pH 6.6-7.3), slightly alkaline (pH 7.4-7.8), moderately alkaline (pH 7.9-8.4), strongly alkaline (pH 8.5-9.0) according to Soil Survey Manual (IARI, 1970). The soils are rated as low (below 0.50 %), medium (0.50-0.75 %) and high (above 0.75 %) in case of organic carbon, low (<280  $\text{kg ha}^{-1}$ ), medium (280 to 560  $\text{kg ha}^{-1}$ ) and high (>560  $\text{kg ha}^{-1}$ ) in case of available nitrogen, low (< 10  $\text{kg ha}^{-1}$ ), medium (10 to 25  $\text{kg ha}^{-1}$ ) and high (> 25  $\text{kg ha}^{-1}$ ) for available phosphorus, low (< 108  $\text{kg ha}^{-1}$ ), medium (108 to 280  $\text{kg ha}^{-1}$ ) and high (> 280  $\text{kg ha}^{-1}$ ) for available potassium and low (<10  $\text{mg kg}^{-1}$ ), medium (10-20  $\text{mg kg}^{-1}$ ) and high (> 20  $\text{mg kg}^{-1}$ ) for available sulphur (Singh *et. al.* 2004, Mehta *et. al.*1988). Critical limits of Fe, Mn, Zn, Cu and B, which separate deficient from non-deficient soils followed in India are 4.5, 2.0, 0.5, 0.2 and 0.5  $\text{mg kg}^{-1}$  respectively. (Follet and Lindsay, 1970 and Berger and Truog, 1940).

The maps for the above mentioned parameters have been prepared using Geographic Information System (GIS) from data generated by analysis of grid soil samples.

## 4. SOIL ACIDITY AND FERTILITY STATUS

### 4.1 Soil Reaction

Soil pH is an important soil property, which affects the availability of several plant nutrients. It is a measure of acidity and alkalinity and reflects the status of base saturation. The soils of the district have been grouped under four soil reaction classes according to Soil Survey Manual (IARI, 1970).

All the soils are acidic in reaction. The soil pH ranges from 4.4 to 6.4. The soil reaction classes with area are given in table 2 and figure 2. The data reveals that strongly acidic soils cover 37.6 % area of the district followed by very strongly acidic (35.6 % of TGA), moderately acid (20.7 % of TGA) and slightly acid (5.1 % of TGA).

**Table 2. Soils under different reaction classes**

Soil reaction	Area ('00 ha)	% of the TGA
Very strongly acidic (pH 4.5 to 5.0)	1338	35.6
Strongly acidic (pH 5.1 to 5.5)	1412	37.6
Moderately acidic (pH 5.6 to 6.0)	778	20.7
Slightly acidic (pH 6.1 to 6.5)	191	5.1
Miscellaneous	38	1.0
Total	3757	100.0

### 4.2 Organic Carbon

The effect of soil organic matter on soil properties is well recognized. Soil organic matter plays a vital role in supplying plant nutrients, cation exchange capacity, improving soil aggregation and hence water retention and soil biological activity.

The organic carbon content in the district ranges from 0.04 to 1.75 %. They are mapped into three classes i.e., low (below 0.5 %), medium (0.5-0.75 %) and high (above 0.75 %). The details are given in table 3 and figure 3. From table 3 it is seen that 46.2 percent area have low surface organic carbon content.



Medium and high organic carbon content constitute 39.4 and 13.4 percent area respectively. Low organic carbon content may be due to high temperature regime of the area, which accelerates the rate of decomposition of organic carbon.

**Table 3. Organic carbon status**

<b>Organic carbon (%)</b>	<b>Area ('00 ha)</b>	<b>% of the TGA</b>
Low (below 0.50 %)	1737	46.2
Medium (0.50-0.75 %)	1479	39.4
High (above 0.75 %)	503	13.4
Miscellaneous	38	1.0
Total	3757	100.0

### **4.3 Macronutrients**

Nutrients like nitrogen (N), phosphorus (P) and potassium (K) are considered as primary nutrients and sulphur (S) as secondary nutrient. These nutrients help in proper growth, development and yield differentiation of plants and are generally required by plants in large quantity.

#### **4.3.1 Available Nitrogen**

Nitrogen is an integral component of many compounds including chlorophyll and enzyme essential for plant growth. It is an essential constituent for amino acids which is building blocks for plant tissue, cell nuclei and protoplasm. It encourage aboveground vegetative growth and deep green colour to leaves. Deficiency of nitrogen decreases rate and extent of protein synthesis and result into stunted growth and develop chlorosis.

Available nitrogen content in the surface soils of the district ranges between 56 and 561 kg/ha and details are given in table 4 and figure 4. Majority area (56.6 % of TGA) of the district have low availability status of available

nitrogen (Below 280 kg ha<sup>-1</sup>) and 41.6 percent area have medium available nitrogen content (280-560 kg ha<sup>-1</sup>).

**Table 4. Available nitrogen status in the surface soils**

<b>Available nitrogen (kg/ha)</b>	<b>Area ('00 ha)</b>	<b>% of the TGA</b>
Low (below 280)	2126	56.6
Medium (280-560)	1564	41.6
High (above 560)	29	0.8
Miscellaneous	38	1.0
Total	3757	100.0

#### **4.3.2 Available Phosphorus**

Phosphorus is important component of adenosine di-phosphate (ADP) and adenosine tri-phosphate (ATP), which involves in energy transformation in plant. It is essential component of deoxyribonucleic acid (DNA), the seat of genetic inheritance in plant and animal. Phosphorous take part in important functions like photosynthesis, nitrogen fixation, crop maturation, root development, strengthening straw in cereal crops etc. The availability of phosphorous is restricted under acidic and alkaline soil reaction mainly due to P-fixation. In acidic condition it get fixed with aluminum and iron and in alkaline condition with calcium.

Available phosphorus content in these soils ranges between 1.0 and 25.7 kg/ha and area and distribution is given in table 5 and figure 5. Majority of the soils (90.5 % area) are low in available phosphorous content (below 10 kg ha<sup>-1</sup>). Soils of 8.2 percent area have medium (10-25 kg ha<sup>-1</sup>) and 0.3 percent area have high (above 25 kg ha<sup>-1</sup>) in available phosphorous content.

**Table 5. Available phosphorous status in the surface soils**

<b>Available phosphorous (kg/ha)</b>	<b>Area ('00 ha)</b>	<b>% of the TGA</b>
Low (below 10)	3401	90.5
Medium (10-25)	307	8.2
High (above 25)	11	0.3
Miscellaneous	38	1.0
Total	3757	100.0

**4.3.3 Available Potassium**

Potassium is an activator of various enzymes responsible for plant processes like energy metabolism, starch synthesis, nitrate reduction and sugar degradation. It is extremely mobile in plant and help to regulate opening and closing of stomata in the leaves and uptake of water by root cells. It is important in grain formation and tuber development and encourages crop resistance for certain fungal and bacterial diseases.

Available potassium content in these soils ranges between 45 and 1019 kg/ha and details about area and distribution is given in table 6 and figure 6. The data reveals that 49 % area of the district have high available potassium content (above 280 kg ha<sup>-1</sup>). Soils of 43.9 percent area are medium (108-280 kg ha<sup>-1</sup>) and 6.1 percent area are low (below 108 kg ha<sup>-1</sup>) in available potassium content.

**Table 6. Available potassium status in the surface soils**

<b>Available potassium (kg/ha)</b>	<b>Area ('00 ha)</b>	<b>% of the TGA</b>
Low (below 108)	230	6.1
Medium (108-280)	1648	43.9
High (above 280)	1841	49.0
Miscellaneous	38	1.0
Total	3757	100.0

#### 4.3.4 Available Sulphur

Sulphur is essential in synthesis of sulphur containing amino acids (cystine, cysteine and methionine), chlorophyll and metabolites including co-enzyme A, biotin, thiamine, or vitamin B1 and glutathione. It activates many proteolytic enzymes, increase root growth & nodule formation and stimulate seed formation.

The available sulphur content in the soils ranges from 2.33 to 51.79 mg kg<sup>-1</sup> and details about area and distribution is given in table 7 and figure 7. Soils of 25.8 percent of the area are low (<10mg kg<sup>-1</sup>) whereas soils of 42.3 and 30.9 percent area are medium and high in available sulphur content.

**Table 7. Available sulphur status in the surface soils**

<b>Available sulphur (mg kg<sup>-1</sup>)</b>	<b>Area ('00 ha)</b>	<b>% of the TGA</b>
Low (<10)	968	25.8
Medium (10-20)	1588	42.3
High (>20)	1163	30.9
Miscellaneous	38	1.0
Total	3757	100.0

#### 4.4 Micronutrients

Proper understanding of micronutrients availability in soils and extent of their deficiencies is the pre-requisite for efficient management of micronutrient fertilizer to sustain crop productivity. Therefore, it is essential to know the micronutrients status of soil before introducing any type of land use.

##### 4.4.1 Available Iron

Iron is constituent of cytochromes, haems and nonhaem enzymes. It is capable of acting as electron carrier in many enzyme systems that bring about oxidation-reduction reactions in plants. It promotes starch formation and seed maturation.

The available iron content in the surface soils is ranges between 9.2 and 552.0 mg kg<sup>-1</sup>. As per the critical limit of available iron (> 4.5 mg kg<sup>-1</sup>), all the soils are sufficient in available iron. They are grouped and mapped into six classes. Majority of the soils (52.3 % of TGA) have available iron content between the range of 50 to 200 mg kg<sup>-1</sup>. The details of area and distribution is presented in table 8 and figure 8.

**Table 8. Available iron status in the surface soils**

<b>Available iron (mg kg<sup>-1</sup>)</b>	<b>Area ('00 ha)</b>	<b>% of the TGA</b>	<b>Rating</b>
<15	237	6.3	Sufficient
15-25	279	7.4	
25-50	611	16.3	
50-100	1013	27.0	
100-200	951	25.3	
200-400	628	16.7	
Miscellaneous	38	1.0	
Total	3757	100.0	

#### **4.4.2 Available Manganese**

Manganese is essential in photosynthesis and nitrogen transformations in plants. It activates decarboxylase, dehydrogenase, and oxidase enzymes.

The available manganese content in surface soils ranges between 2.4 and 498.0 mg kg<sup>-1</sup>. As per the critical limit of available manganese (> 2 mg kg<sup>-1</sup>), all the soils are sufficient in available manganese. They are grouped and mapped into six classes. Majority of soils 71.1 % of TGA of district have available Mn content between 50 to 200 mg kg<sup>-1</sup>. The details of area and distribution are presented in table 9 and figure 9.

**Table 9. Available manganese status in the surface soils**

Available manganese (mg kg <sup>-1</sup> )	Area ('00 ha)	% of the TGA	Rating
<10	95	2.5	Sufficient
10-25	120	3.2	
25-50	596	15.9	
50-100	1447	38.5	
100-200	1224	32.6	
200-500	237	6.3	
Miscellaneous	38	1.0	
Total	3757	100.0	

**4.4.3 Available Zinc**

Zinc plays role in protein synthesis, reproductive process of certain plants and in the formation starch and some growth hormones. It promotes seed maturation and production.

The available zinc in surface soils ranges between 0.45 and 2.24 mg kg<sup>-1</sup>. They are grouped and mapped into four classes. Majority of area (98.6 % of TGA) have sufficient amount of available zinc (> 0.5 mg kg<sup>-1</sup>) and soils of 0.4 per cent area are deficient (< 0.5 mg kg<sup>-1</sup>) in available zinc. The details of area and distribution are presented in table 10 and figure 10.

**Table 10. Available zinc status in the surface soils**

Available zinc (mg kg <sup>-1</sup> )	Area ('00 ha)	% of the TGA	Rating
<0.5	14	0.4	Deficient
0.5-1.0	230	6.1	Sufficient
1.0-2.0	269	7.2	
2.0-3.0	3206	85.3	
Miscellaneous	38	1.0	
Total	3757	100.0	

#### 4.4.4 Available Copper

Copper involves in photosynthesis, respiration, protein and carbohydrate metabolism and in the use of iron. It stimulates lignifications of all the plant cell wall and is capable of acting as electron carrier in many enzyme systems that bring about oxidation-reduction reactions in plants.

The available copper status in surface soils ranges between 0.15 and 3.18 mg kg<sup>-1</sup>. Majority of soils (96.7 % of TGA) have sufficient amount of available copper (>0.2 mg kg<sup>-1</sup>) and soils of 2.3 % area are deficient in available copper (<0.2 mg kg<sup>-1</sup>). They are grouped and mapped into six classes. The details of area and distribution are presented in table 11 and figure 11.

**Table 11. Available copper status in the surface soils**

<b>Available copper (mg kg<sup>-1</sup>)</b>	<b>Area (‘00 ha)</b>	<b>% of the TGA</b>	<b>Rating</b>
<0.2	88	2.3	Deficient
0.2-0.5	473	12.6	Sufficient
0.5-1.0	1141	30.4	
1.0-2.0	1581	42.1	
2.0-4.0	436	11.6	
Miscellaneous	38	1.0	
Total	3757	100.0	

#### 4.4.5 Available Boron

Boron increases solubility and mobility of calcium in the plant and it act as regulator of K/Ca ratio in the plant. It is required for development of new meristematic tissue and also necessary for proper pollination, fruit and seed setting and translocation of sugar, starch and phosphorous etc. It has role in synthesis of amino acid and protein and regulates carbohydrate metabolism.

The available boron content in the soils ranges from 0.01 to 2.28 mg kg<sup>-1</sup> and details about area and distribution is given in table 12 and figure 12. The critical limit for deficiency of the available boron is <0.5. Soils of 45.7 percent

area of district are deficient ( $<0.50 \text{ mgkg}^{-1}$ ) whereas 53.3 percent area have sufficient ( $>0.50 \text{ mgkg}^{-1}$ ) in available boron content.

**Table 12. Available boron status in the surface soils**

<b>Available boron (<math>\text{mg kg}^{-1}</math>)</b>	<b>Area (<math>'00 \text{ ha}</math>)</b>	<b>% of the TGA</b>	<b>Rating</b>
<0.25	861	22.9	Deficient
0.25-0.50	857	22.8	
0.50-0.75	765	20.4	Sufficient
>0.75	1236	32.9	
Miscellaneous	38	1.0	
Total	3757	100.0	



## 5. SUMMARY

All the soils are acidic in reaction and pH ranges from 4.4 to 6.4. Strongly acidic soils cover 37.6 % area of the district followed by very strongly acidic (35.6 % of TGA), moderately acid (20.7 % of TGA) and slightly acid (5.1 % of TGA). The organic carbon content in the district ranges from 0.04 to 1.75 %. Soils of 46.2 percent area have low surface organic carbon content. Medium and high organic carbon content constitute 39.4 and 13.4 percent area respectively.

Available nitrogen content in the surface soils of the district ranges between 56 and 561 kg/ha. Majority area (56.6 % of TGA) of the district have low availability status of available nitrogen (Below 280 kg ha<sup>-1</sup>) and 41.6 percent area have medium available nitrogen content (280-560 kg ha<sup>-1</sup>). Available phosphorus content in these soils ranges between 1.0 and 24.7 kg/ha. Majority of the soils (90.5 % area) are low in available phosphorous content (below 10 kg ha<sup>-1</sup>). Available potassium content in these soils ranges between 45 and 1019 kg/ha. The data reveals that 49.0 % area of the district have high available potassium content (above 280 kg ha<sup>-1</sup>). Soils of 43.9 percent area are medium (108-280 kg ha<sup>-1</sup>) and 6.1 percent area are low (below 108 kg ha<sup>-1</sup>) in available potassium content. Available sulphur content ranges between 2.33 to 51.79 mg kg<sup>-1</sup>. Soils of 25.8 percent of the area are low (<10 mg kg<sup>-1</sup>) whereas soils of 42.3 and 30.9 percent area are medium and high in available sulphur content.

All the soils of district are sufficient in available iron and manganese whereas soils of 0.4 and 2.3 percent area are deficient in available zinc and copper respectively. Available boron content in the soils ranges between 0.01 and 2.28 mg kg<sup>-1</sup> and 45.7 percent area of district are deficient (<0.50 mg kg<sup>-1</sup>).

## REFERENCES

- Berger, K. C. and Truog, E.(1940) J. Am. Soc. Agron. 32,297
- Census of India (2001) Primary Census Abstract.
- FAI (2003 – 2004) Fertiliser and agriculture statistics, Eastern Region.
- Follet, R. H. and Lindsay , W. L.(1970) Tech. Bull. Colo. Agric. Exp. Station 110.
- Haldar, A. K., Srivastava, R., Thampi, C. J., Sarkar, D., Singh, D. S., Sehgal, J and Velayutham, M. (1996) Soils of Bihar for optimizing land use. **NBSS Publ. 50b.** (Soils of India Series), National Bureau of Soil Survey and Land Use Planning, Nagpur, India, pp. 70+4 sheets soil Map (1:500,000 scale).
- Hatcher, J. T. and Wilcox, L. V. (1950) Analyt. Chem. **22**, 567
- I.A.R.I. (1970). Soil survey manual, All India Soil and Land Use Organization, Indian Agricultural Research Institute, New Delhi.
- Lindsay , W. L. & Norvell, W.A.(1978). Development of a DTPA micronutrients soil test for Zn, Fe, Mn and Cu. *Soil Sci. Soc. Am. Proceedings.* 42, 421-428
- Mehta, V. S. , Singh, V and Singh, R. P. (1988) J. Indian Society of Soil Science, 36, 743
- Mishra, R. K. (2004) Planning for Food and Nutritional Security in Jharkhand, *Published by Agricultural Data Bank, BAU, Ranchi, Jharkhand*, p. 275
- Page, A. L., Miller, R. H. and Keeney, D. R.(1982) Method of Soil Analysis, Part-II, Chemical and Microbiological Properties, Soil Sci. Soc. Am. And Am. Soc. Agron. Madison, Wisconsin, USA.
- Singh Dhyan, Chhonkar, P. K. and Pandey, R. N.(2004) Soil Plant and Water Analysis, A Manual, IARI, New Delhi.
- Subbaiah, B.V. and Asija, G.L. (1956). A rapid procedure for determination of available nitrogen in soil. *Current Science* 25, 259-260
- Tandon, H. L. S. (Ed) (1999) *Methods of analysis of soils, plants, waters and fertilizers.* Fertilizer Development and Consultation Organisation, New Delhi, India.
- William, C. H. and Stainbergs, A. (1959) Aust. J. Agric. Res. **10**, 342