

**ASSESSMENT AND MAPPING OF SOME IMPORTANT SOIL
PARAMETERS INCLUDING SOIL ACIDITY FOR THE STATE OF
JHARKHAND (1:50,000 SCALE) TOWARDS
RATIONAL LAND USE PLAN**

JAMTARA DISTRICT



*National Bureau of Soil Survey and Land Use Planning (ICAR)
Regional Centre, Kolkata*

*In collaboration with :
Deptt. Of Soil Science & Agricultural Chemistry, BAU, Ranchi, Jharkhand*

*Sponsored by : Department of Agriculture & Cane Development,
Govt. of Jharkhand*

1. INTRODUCTION

Reliable information on the location, extent and quality of soil and land resources is the first requirement in planning for the sustainable management of land resources. The components of land i.e., soils, climate, water, nutrient and biota are organised into eco-system which provide a variety of services that are essential to the maintenance of the life support system and the productive capacity of the environment. Our land mass is fixed, but the competition among different kinds of uses for this land is increasing because of rapidly rising global population. Therefore, integrated land resource planning and management are required to resolve these conflicts and soil resource survey seems to be a viable means in this process and knowledge of soil fertility status and problems of soils like soil acidity/alkalinity become essential for sustainable land use plan.

Soil fertility is an aspect of the soil-plant relationship. Fertility status of the soils is primarily and importantly dependent upon both the macro and micronutrient reserve of that soil. Continued removal of nutrients by crops, with little or no replacement will increase the nutrient stress in plants and ultimately lowers the productivity. The fertility status of the soils mainly depends on the nature of vegetation, climate, topography, texture of soil and decomposition rate of organic matter. Optimum productivity of any cropping systems depends on adequate supply of plant nutrients. GIS is a versatile tool used for integration of soil database and production of a variety of users specific and user-friendly interpretative maps. This further leads to accurately and scientifically interpret and plan some of the aspects like conservation of organic matter, soil reaction (pH) control and fertilization.

Keeping in view NBSS & LUP, Regional Centre, Kolkata in collaboration with Department of Soil Science and Agricultural Chemistry, BAU, Ranchi, Jharkhand undertook a project entitled "Assessment and mapping of some important soil parameters including soil acidity for the state of Jharkhand (1:50,000 scale)

towards rational land use plan” from Department of Agriculture, Govt. of Jharkhand. The major objectives of the project were

- Preparation of districtwise soil acidity maps
- Preparation of districtwise soil fertility maps (Organic carbon, available N, P, K, S and available Fe, Mn, Zn, Cu and B)

The above maps will provide information regarding soil nutrients and soil acidity status for the districts, which will be very useful in identification of site specific problems for planning purposes. The present report deals with the above mentioned objectives of the Jamtara district, Jharkhand.

2. GENERAL DESCRIPTION OF THE AREA

2.1 Location and Extent

Jamtara is new district created by separating from Dumka district. It is bounded by Deoghar district in north, Dumka and West Bengal in the east, Dhanbad and West Bengal in the south and Giridih in the west. It has an area of 1802 sq. km and population of 6,97,059 persons (Census of India, 2001). The district comprises one subdivision (Jamtara) and four development blocks viz. Kundahit, Nala, Jamtara and Narayanpur.

2.2 Physiography, Geology and Drainage

The district is located in Rajmahal highland. It consist of rolling open land surface with long ridges and intervening depressions. Average elevation of the district ranges between 150 to 300 metres. Geologically the area is comprised with basaltic trap and sedimentary beds. Quartz and gneiss are found at some places. Ajay is the major river flowing through the district.

2.3 Climate

Owing to its position near West Bengal and hilly landscape of the region climatic condition is slightly different from the rest of the state. The district receives an annual rainfall of 1500 mm. and most of the rainfall occurs during the rainy season. During winter season the temperature varies between 16 to 21⁰ C and during summer season it varies between 22 to 36⁰ C.

2.4 Agriculture and Land Use

Since the area consist of hilly landscape and receive more rainfall it has considerable vegetative cover. But due to ruthless exploitation, most of the forest have turned into bushes devoid of big trees. Important trees include sal, semal, kaha, ashwar, mahua etc. important crops of the district are paddy, maize

and wheat. Only 6.42 percent area of agricultural use is net irrigated and major source of irrigations are well and tanks.

Land Use in Jamtara District (1997-98)

	Jamtara	Jharkhand
1. Forest	11.27 %	29.2 %
2. Net sown area	29.47 %	22.7 %
3. Barren and unculturable waste	6.08 %	7.2 %
4. Non agricultural use	12.38 %	9.9 %
5. Orchards	1.77 %	2.5 %
6. Pasture	4.90 %	
7. Culturable wasteland	6.18 %	3.5 %
8. Current and other fallow	27.95 %	25.0 %

Source: Fertilizer and Agriculture Statistics, Eastern Region (2003-2004)

2.5 Soils

The soils occurring in different landforms have been characterised during soil resource mapping of the state on 1:250,000 scale (Haldar *et al.* 1996) and three soil orders namely Entisols, Inceptisols and Alfisols were observed in Jamtara district (Fig.1 and table 1). Inceptisols were the dominant soils covering 50.8 percent of TGA followed by Alfisols (39.3 %) and Entisols (6.6 %).

Table 1. Soils of the district and their extent

Map unit	Taxonomy	Area ('00ha)	% of TGA
23	Fine-loamy, mixed, hyperthermic, Typic Haplustepts Fine-loamy, mixed, hyperthermic, Typic Haplustalfs	70	3.88
25	Fine, mixed, hyperthermic Typic Paleustalfs Fine, mixed, hyperthermic Rhodic Paleustalfs	24	1.33
29	Loamy, mixed, hyperthermic Lithic Haplustepts Fine-loamy, mixed, hyperthermic Typic Ustorthents	294	16.32
30	Loamy-skeletal, mixed, Typic Haplustepts Fine-loamy, mixed, hyperthermic Typic Haplustalfs	278	15.43
31	Fine-loamy, mixed, hyperthermic Typic Haplustepts Fine, mixed, hyperthermic Typic Paleustalfs	814	45.17
32	Fine-loamy, mixed, hyperthermic Typic Haplustepts Coarse loamy, mixed, hyperthermic Typic Ustorthents	2	0.11
34	Fine loamy, mixed, hyperthermic Typic Paleustalfs Fine-loamy, mixed, hyperthermic Typic Rhodustalfs	220	12.21
44	Fine, mixed, hyperthermic Aeric Endoaquepts Fine, mixed, hyperthermic Typic Haplustepts	41	2.28
Miscellaneous		59	3.27
Total		1802	100.00

3. METHODOLOGY

The base map of the district was prepared on 1:50,000 scale using Survey of India toposheets (72L/8,12,16,72P/4,73I/9,13 and 73M/1,5) and all the maps were demarcated with grid points at 2.5 km interval.

Surface soil samples from demarcated grid points and other related informations were collected through field survey. Soil samples were air dried, processed and analysed for pH, organic carbon, available phosphorous and potassium (Page *et al.*, 1982), available nitrogen (Subbaiah and Asija, 1956), available sulphur by using 0.15 percent CaCl_2 as the extractant (William and Steinbergs, 1959), available (DTPA extractable) Fe, Mn, Zn and Cu (Lindsay and Norvell, 1978) and available B (hot water soluble) by Carmine method (Hatcher and Wilcox, 1950).

The soils are grouped under different soil reaction classes viz extremely acidic (pH<4.5), very strongly acidic (pH 4.5 – 5.0), strongly acidic (pH 5.1 – 5.5), moderately acidic (pH 5.6-6.0), slightly acidic (pH 6.1-6.5), neutral (pH 6.6-7.3), slightly alkaline (pH 7.4-7.8), moderately alkaline (pH 7.9-8.4), strongly alkaline (pH 8.5-9.0) according to Soil Survey Manual (IARI, 1970).The soils are rated as low (below 0.50 %), medium (0.50-0.75 %) and high (above 0.75 %) in case of organic carbon, low (<280 kg ha⁻¹), medium (280 to 560 kg ha⁻¹) and high (>560 kg ha⁻¹) in case of available nitrogen, low (< 10 kg ha⁻¹), medium (10 to 25 kg ha⁻¹) and high (> 25 kg ha⁻¹) for available phosphorus, low (< 108 kg ha⁻¹), medium (108 to 280 kg ha⁻¹) and high (> 280 kg ha⁻¹) for available potassium and low (<10 mg kg⁻¹), medium (10-20 mg kg⁻¹) and high (> 20 mg kg⁻¹) for available sulphur (Singh *et. al.* 2004, Mehta *et. al.*1988). Critical limits of Fe, Mn, Zn, Cu and B, which separate deficient from non-deficient soils followed in India are 4.5, 2.0, 0.5, 0.2 and 0.5 mg kg⁻¹ respectively. (Follet and Lindsay, 1970 and Berger and Truog, 1940).

The maps for the above mentioned parameters have been prepared using Geographic Information System (GIS) from data generated by analysis of grid soil samples.

4. SOIL ACIDITY AND FERTILITY STATUS

4.1 Soil Reaction

Soil pH is an important soil property, which affects the availability of several plant nutrients. It is a measure of acidity and alkalinity and reflects the status of base saturation. The soils of the district have been grouped under seven soil reaction classes according to Soil Survey Manual (IARI, 1970).

The soil pH ranges from 4.3 to 7.4. The soil reaction classes with area are given in table 2 and figure 2. The data reveals that majority of the area is acidic (88 % of TGA), in which 36.6 percent area is strongly acidic, 25.6 percent very strongly acidic, 18.1 percent moderately acidic, 5.6 percent slightly acidic and 2.1 percent extremely acidic in reaction. Soils of 5.4 percent area of the district are neutral whereas 3.3 percent area is slightly alkaline in reaction.

Table 2. Soils under different reaction classes

Soil reaction	Area ('00ha)	% of the TGA
Extremely acidic (pH 4.0 to 4.5)	37	2.1
Very strongly acidic (pH 4.6 to 5.0)	462	25.6
Strongly acidic (pH 5.1 to 5.5)	659	36.6
Moderately acidic (pH 5.6 to 6.0)	327	18.1
Slightly acidic (pH 6.1 to 6.5)	101	5.6
Neutral (pH 6.6 to 7.3)	97	5.4
Slightly alkaline (pH 7.4 to 7.8)	60	3.3
Miscellaneous	59	3.3
Total	1802	100.0

4.2 Organic Carbon

The effect of soil organic matter on soil properties is well recognized. Soil organic matter plays a vital role in supplying plant nutrients, cation exchange capacity, improving soil aggregation and hence water retention and soil biological activity.

The organic carbon content in the district ranges from 0.16 to 1.47 %. They are mapped into three classes i.e., low (below 0.5 %), medium (0.5-0.75 %) and high (above 0.75 %) (Table 3 and Figure 3). From table 3 it is seen that 47.4 percent area of the district shows high organic carbon content. Medium and low organic carbon content constitute 24.7 and 24.6 percent area respectively.

Table 3. Organic carbon status

Organic carbon (%)	Area ('00ha)	% of the TGA
Low (below 0.50 %)	444	24.6
Medium (0.50-0.75 %)	445	24.7
High (above 0.75 %)	854	47.4
Miscellaneous	59	3.3
Total	1802	100.0

4.3 Macro Nutrients

Nutrients like nitrogen (N), phosphorus (P) and potassium (K) are considered as primary nutrients and sulphur (S) as secondary nutrient. These nutrients help in proper growth, development and yield differentiation of plants and are generally required by plants in large quantity.

4.3.1 Available Nitrogen

Nitrogen is an integral component of many compounds including chlorophyll and enzyme essential for plant growth. It is an essential constituent for amino acids which is building blocks for plant tissue, cell nuclei and protoplasm. It encourage aboveground vegetative growth and deep green colour to leaves. Deficiency of nitrogen decreases rate and extent of protein synthesis and result into stunted growth and develop chlorosis.

Available nitrogen content in the surface soils of the district ranges between 135 and 543 kg/ha and details are given in table 4 and figure 4. Majority soils (76.5 % of TGA) of the district have medium availability status of

nitrogen (280-560 kg ha⁻¹) and soils of 20.2 percent area have low available nitrogen content (<280 kg ha⁻¹).

Table 4. Available nitrogen status in the surface soils

Available nitrogen (kg ha⁻¹)	Area ('00ha)	% of the TGA
Low (below 280)	365	20.2
Medium (280-560)	1378	76.5
Miscellaneous	59	3.3
Total	1802	100.0

4.3.2 Available Phosphorus

Phosphorus is important component of adenosine di-phosphate (ADP) and adenosine tri-phosphate (ATP), which involves in energy transformation in plant. It is essential component of deoxyribonucleic acid (DNA), the seat of genetic inheritance in plant and animal. Phosphorous take part in important functions like photosynthesis, nitrogen fixation, crop maturation, root development, strengthening straw in cereal crops etc. The availability of phosphorous is restricted under acidic and alkaline soil reaction mainly due to P-fixation. In acidic condition it get fixed with aluminum and iron and in alkaline condition with calcium.

Available phosphorus content in these soils ranges between 1.4 and 18.5 kg/ha and their distribution is given in table 5 and figure 5. Data reveals that majority of the soils are low (73.1 % of TGA) followed by medium (23.6 % of TGA) content of available phosphorous.

Table 5. Available phosphorous status in the surface soils

Available phosphorous (kg ha⁻¹)	Area ('00ha)	% of the TGA
Low (below 10)	1318	73.1
Medium (10-25)	425	23.6
Miscellaneous	59	3.3
Total	1802	100.0

4.3.3 Available Potassium

Potassium is an activator of various enzymes responsible for plant processes like energy metabolism, starch synthesis, nitrate reduction and sugar degradation. It is extremely mobile in plant and help to regulate opening and closing of stomata in the leaves and uptake of water by root cells. It is important in grain formation and tuber development and encourages crop resistance for certain fungal and bacterial diseases.

Available potassium content in these soils ranges between 39 and 582 kg/ha and details about area and distribution is given in table 6 and figure 6. The data reveals that most of the soils (61.3 % of TGA) have medium available potassium content (108-280 kg ha⁻¹). Soils of 14.1 percent area are high (above 280 kg ha⁻¹) and 21.3 percent area are low (below 108) in available potassium content.

Table 6. Available potassium status in the surface soils

Available potassium (kg ha⁻¹)	Area ('00ha)	% of the TGA
Low (below 108)	383	21.3
Medium (108-280)	1105	61.3
High (above 280)	255	14.1
Miscellaneous	59	3.3
Total	1802	100.0

4.3.4 Available Sulphur

Sulphur is essential in synthesis of sulphur containing amino acids (cystine, cysteine and methionine), chlorophyll and metabolites including co-enzyme A, biotin, thiamine, or vitamin B1 and glutathione. It activates many proteolytic enzymes, increase root growth and nodule formation and stimulate seed formation.

The available sulphur content in the soils ranges from 0.76 to 41.49 mg kg⁻¹ and details about area and distribution is given in table 7 and figure 7. Soils

of 25.8 percent of the area are low ($<10 \text{ mg kg}^{-1}$) whereas soils of 39.7 and 31.2 percent area are medium ($10\text{-}20 \text{ mg kg}^{-1}$) and high ($>20 \text{ mg kg}^{-1}$) in available sulphur content respectively.

Table 7. Available sulphur status in the surface soils

Available sulphur (mg kg^{-1})	Area ($'00\text{ha}$)	% of the TGA
Low (<10)	465	25.8
Medium ($10\text{-}20$)	715	39.7
High (>20)	563	31.2
Miscellaneous	59	3.3
Total	1802	100.0

4.4 Micronutrients

Proper understanding of micronutrients availability in soils and extent of their deficiencies is the pre-requisite for efficient management of micronutrient fertilizer to sustain crop productivity. Therefore, it is essential to know the micronutrients status of soil before introducing any type of land use.

4.4.1 Available Iron

Iron is constituent of cytochromes, haems and nonhaem enzymes. It is capable of acting as electron carrier in many enzyme systems that bring about oxidation-reduction reactions in plants. It promotes starch formation and seed maturation.

The available iron content in the surface soils is ranges between 5.9 and 82.0 mg kg^{-1} . As per the critical limit of available iron ($> 4.5 \text{ mg kg}^{-1}$), all the soils are sufficient in available iron. They are grouped and mapped into four classes. Majority of the soils (50.6 % of TGA) have available iron content between the range of 50 to 100 mg kg^{-1} . The details of area and distribution is presented in table 8 and figure 8.

Table 8. Available iron status in the surface soils

Available iron (mg kg⁻¹)	Area (⁰⁰ha)	% of the TGA	Rating
<15	116	6.4	Sufficient
15-25	153	8.5	
25-50	562	31.2	
50-100	912	50.6	
Miscellaneous	59	3.3	
Total	1802	100.0	

4.4.2 Available Manganese

Manganese is essential in photosynthesis and nitrogen transformations in plants. It activates decarboxylase, dehydrogenase, and oxidase enzymes.

The available manganese content in surface soils ranges between 5.9 and 83.2 mg kg⁻¹. As per the critical limit of available manganese (> 2 mg kg⁻¹), all the soils are sufficient in available manganese. They are grouped and mapped into four classes. Most of soils (44.5 %) have available Mn content between 10 and 25 mg kg⁻¹. The details of area and distribution are presented in table 9 and figure 9.

Table 9. Available manganese status in the surface soils

Available manganese (mg kg⁻¹)	Area (⁰⁰ha)	% of the TGA	Rating
<10	150	8.3	Sufficient
10-25	802	44.5	
25-50	273	15.2	
50-100	518	28.7	
Miscellaneous	59	3.3	
Total	1802	100.0	

4.4.3 Available Zinc

Zinc plays role in protein synthesis, reproductive process of certain plants and in the formation starch and some growth hormones. It promotes seed maturation and production.

The available zinc in surface soils ranges between 0.38 and 5.00 mg kg⁻¹. They are grouped and mapped into five classes. Majority of soils (92.9 % of TGA) are sufficient (>0.5 mg kg⁻¹) whereas soils of 3.8 percent area are deficient (<0.5 mg kg⁻¹) in available zinc. The details of area and distribution are presented in table 10 and figure 10.

Table 10. Available zinc status in the surface soils

Available zinc (mg kg ⁻¹)	Area ('00ha)	% of the TGA	Rating
<0.5	68	3.8	Deficient
0.5-1.0	303	16.8	Sufficient
1.0-2.0	1011	56.1	
2.0-3.0	253	14.0	
3.0-5.0	108	6.0	
Miscellaneous	59	3.3	
Total	1802	100.0	

4.4.4 Available Copper

Copper involves in photosynthesis, respiration, protein and carbohydrate metabolism and in the use of iron. It stimulates lignifications of all the plant cell wall and is capable of acting as electron carrier in many enzyme systems that bring about oxidation-reduction reactions in plants.

The available copper status in surface soils ranges between 0.14 and 5.06 mg kg⁻¹. They are grouped and mapped into six classes. Majority of soils (95.8 % of TGA) have sufficient amount of available copper (>0.2 mg kg⁻¹) and soils of 0.9 % area are deficient in available copper (<0.2 mg kg⁻¹). The details of area and distribution are presented in table 11 and figure 11.

Table 11. Available copper status in the surface soils

Available copper (mg kg ⁻¹)	Area ('00ha)	% of the TGA	Rating
<0.2	16	0.9	Deficient
0.2-0.5	50	2.8	Sufficient
0.5-1.0	58	3.2	
1.0-2.0	290	16.1	
2.0-4.0	1208	67.0	
4.0-6.0	121	6.7	
Miscellaneous	59	3.3	
Total	1802	100.0	

4.4.5 Available Boron

Boron increases solubility and mobility of calcium in the plant and it act as regulator of K/Ca ratio in the plant. It is required for development of new meristematic tissue and also necessary for proper pollination, fruit and seed setting and translocation of sugar, starch and phosphorous etc. It has role in synthesis of amino acid and protein and regulates carbohydrate metabolism.

The available boron content in the soils ranges from 0.02 to 6.09 mgkg⁻¹ and details about area and distribution is given in table 12 and figure 12. The critical limit for deficiency of the available boron is <0.5. Soils of 23.0 percent area of district are deficient (<0.50 mgkg⁻¹) whereas 73.7 percent area are sufficient (>0.50 mgkg⁻¹) in available boron content.

Table 12. Available boron status in the surface soils

Available boron (mg kg ⁻¹)	Area ('00ha)	% of the TGA	Rating
<0.25	290	16.1	Deficient
0.25-0.50	125	6.9	
0.50-0.75	169	9.4	Sufficient
>0.75	1159	64.3	
Miscellaneous	59	3.3	
Total	1802	100.0	

5. SUMMARY

The soil pH ranges from 4.3 to 7.4. Soils of 88 % area are acidic in reaction, in which 36.6 percent area strongly acidic, 25.6 percent very strongly acidic, 18.1 percent moderately acidic, 5.6 percent slightly acidic and 2.1 percent extremely acidic in reaction. Soils of 5.4 percent area of the district are neutral whereas 3.3 percent area is slightly alkaline in reaction. The organic carbon content in the district ranges from 0.16 to 1.47 %. Soils of 47.4 percent area of the district shows high organic carbon content. Medium and low organic carbon content constitute 24.7 and 24.6 percent area respectively.

Available nitrogen content in the surface soils of the district ranges between 135 and 543 kg/ha. Majority soils of the district have medium availability of nitrogen and soils of 20.2 percent area have low available nitrogen content. Available phosphorus content in these soils ranges between 1.4 and 18.5 kg/ha. Majority of the soils are low (73.1 % of TGA) followed by medium (23.6 % of TGA) in of available phosphorous content. Available potassium content in these soils ranges between 39 and 582 kg/ha. Most of the soils (61.3 % of TGA) have medium available potassium content. Soils of 14.1 percent area have high and 21.3 percent area have low in available potassium content. The available sulphur content in the soils ranges from 0.76 to 41.49 mg kg⁻¹. Soils of 25.8 percent of the area are low (<10 mg kg⁻¹) whereas soils of 39.7 and 31.2 percent area are medium (10-20 mg kg⁻¹) and high (>20 mg kg⁻¹) in available sulphur content respectively.

Soils are analysed for available (DTPA extractable) micronutrients and seen that all the soils are sufficient in available iron and manganese whereas soils of 3.8 and 0.9 percent area are deficient in available zinc and copper respectively. Available boron content in the soils ranges between 0.02 to 6.09 mgkg⁻¹ and 23.0 percent area of district is deficient (<0.50 mg kg⁻¹) in it.

REFERENCES

- Berger, K. C. and Truog, E.(1940) J. Am. Soc. Agron. 32,297
- Census of India (2001) Primary Census Abstract.
- FAI (2003 – 2004) Fertiliser and agriculture statistics, Eastern Region.
- Follet, R. H. and Lindsay , W. L.(1970) Tech. Bull. Colo. Agric. Exp. Station 110.
- Haldar, A. K., Srivastava, R., Thampi, C. J., Sarkar, D., Singh, D. S., Sehgal, J and Velayutham, M. (1996) Soils of Bihar for optimizing land use. **NBSS Publ. 50b.** (Soils of India Series), National Bureau of Soil Survey and Land Use Planning, Nagpur, India, pp. 70+4 sheets soil Map (1:500,000 scale).
- Hatcher, J. T. and Wilcox, L. V. (1950) Analyt. Chem. **22**, 567
- I.A.R.I. (1970). Soil survey manual, All India Soil and Land Use Organization, Indian Agricultural Research Institute, New Delhi.
- Lindsay , W. L. & Norvell, W.A.(1978). Development of a DTPA micronutrients soil test for Zn, Fe, Mn and Cu. *Soil Sci. Soc. Am. Proceedings.* 42, 421-428
- Mehta, V. S. , Singh, V and Singh, R. P. (1988) J. Indian Society of Soil Science, 36, 743
- Mishra, R. K. (2004) Planning for Food and Nutritional Security in Jharkhand, *Published by Agricultural Data Bank, BAU, Ranchi, Jharkhand*, p. 275
- Page, A. L., Miller, R. H. and Keeney, D. R.(1982) Method of Soil Analysis, Part-II, Chemical and Microbiological Properties, Soil Sci. Soc. Am. And Am. Soc. Agron. Madison, Wisconsin, USA.
- Singh Dhyani, Chhonkar, P. K. and Pandey, R. N.(2004) Soil Plant and Water Analysis, A Manual, IARI, New Delhi.
- Subbaiah, B.V. and Asija, G.L. (1956). A rapid procedure for determination of available nitrogen in soil. *Current Science* 25, 259-260
- Tandon, H. L. S. (Ed) (1999) *Methods of analysis of soils, plants, waters and fertilizers.* Fertilizer Development and Consultation Organisation, New Delhi, India.
- William, C. H. and Stainbergs, A. (1959) Aust. J. Agric. Res. **10**, 342